EFFECT OF THE PROBIOTIC BIOSTIM LBS IN ACUTE INTOXICATION AFTER INDUSTRIAL ACCIDENTS

Georgieva M.¹, N. Alexandrov²

¹Department of Preclinical and Clinical Pharmacology and Toxicology,
Medical University "Prof. Paraskev Stoyanov" - Varna,

²Department of Toxicology and Alergology, Military Medical Academy of Sofia

Reviewed by: assoc. prof. B. Galunska

ABSTRACT

A probiotic is a "live microbial food ingredient that, when ingested in sufficient quantities, exerts health benefits on the consumer". The profound research of a range of Bulgarian scientists has allowed for the development in the last 15-20 years of original Bulgarian probiotic, milk-acid low-lactose products, united in the Biosim LBS (Lactoflor, Biomilk) series. This survey gives an overview of the effect of probiotic Biostim LBS in acute intoxications after industrial accidents. The mean energy values in kcal of the nutritional therapy applied intravenously and via oral route in 186 patients with intoxications with moderate severity are presented. The data show that Biostim LBS covers a mean energy value of 45.36% of the food applied orally in the group with moderate and 70.86% in the group of patients with severe intoxication. These results have been achieved owing to the correct choice of therapy which have included mutually additive enteral and parenteral medications. In none of the patients adverse events have been observed regarding the parenteral medications and the enterally introduced Biostim LBS as well. They exerted beneficial effect on the metabolism, improved body energy levels and contributed to the absorption of energy and proteins. A special attention should be paid to the probiotic properties of the viable cells of *L. bulgaricus*, which favorably influence on the organism of the patients by improvement and regulation of the internal bacterial equilibrium.

Key words: probiotic Biostim LBS, *Lactobacillus bulgaricus*, intoxication, energy, nutritional therapy

INTRODUCTION

A probiotic is a "live microbial food ingredient that, when ingested in sufficient quantities, exerts health benefits on the consumer". Probiotics exert their benefits through several mechanisms. They prevent colonization, cellular adhesion and invasion by pathogenic organisms. Besides they have a direct antimicrobial activity and modulate the host immune response (2,4,8-10). The strongest evidence of the clinical effectiveness of probiotics is provided by their use for the treatment of acute diarrhea, attenuation of antibiotic-associated gastrointestinal side effects, and prevention and treatment of allergy manifestations (5-7,11-13,16).

The profound research of a range of Bulgarian scientists has allowed for the development in the last 15-20 years of original Bulgarian probiotic, milk-acid low-lactose products united in the Biosim (Lactoflor, Biomilk) series. Their composition is represented on Table 1 (1).

The aim of this survey was to give an overview of the effect of probiotic Biomilk in acute intoxications after industrial accidents.

MATERIAL AND METHODS

The dairy product Biomilk was applied as nutrition therapy agent in combination with parenteral solutions. The study covered 234 adult patients of both genders presenting with different types of acute intoxication. Of them, 48 patients presented with severe intoxications and 186 ones - with moderate intoxications. They were hospitalized in the Clinic of Toxicology, Naval Hospital of Varna, in the Clinic of Toxicology at the Military Medical Academy and N.I.Pirogov of Sofia. The intoxications resulted from a toxic smoke during fire; inhalatory poisoning with gasoline in a closed room; as a consequence of cleaning of tanks containing petroleum oil products and of industrial accidents with polyvinyl chloride and dichlorethane.

The therapy schedule included enteral and parenteral medications which ensured the necessary energy for 24 hours (in kcal) after Harris-Benedict equation. The part of the energy provided by Biomilk peroral intake in doses of 125 g/day in water solu-

Address for correspondence:

M. Georgieva, Dept. of Preclinical and Clinical Pharmacology and Toxicology, Medical University of Varna "Prof. Paraskev Stoyanov"
55 Marin Drinov Str., BG-9002 Varna
e-mail: marieta_md@yahoo.com

tion in the patients with moderate intoxication and 250 g/day in the patients with severe intoxication was estimated. The clinical investigations were timely approved by the Ethical Commission of the Higher Military Medical Institute at the Military Medical Academy of Sofia, and The National Emergency Institute "N. Pirogov" of Sofia (Protocol No. 23/10.XII.1986).

RESULTS

The mean energy values in kcal, of the nutritional therapy applied intravenously and via oral route in 186 patients with moderate intoxications are presented on Fig. 1.

Fig. 1. Values of the energy requirements and energy intake in kcal via oral and venous route in patients with moderate intoxications

Fig. 2 demonstrates the mean energy values in kcal applied as a nutritional therapy intravenously and via oral route in 48 patients with severe intoxications.

Fig. 2. Values of the energy requirements and energy intake in kcal via oral and venous route in patients with severe intoxications

Our data show that Biostim LBS covers a mean energy value of 45.36% of the food applied orally in the group of patients with moderate and of 70.86% in the group with severe intoxications.

DISCUSSION

An optimal nutrition should meet not only energy requirements. By means of adequate import of macro-

microelements and vitamins it should ensure the fulfillment of the complex vital processes in the body such as oxidation, degradation, synthesis of new substances and hormones, thermoregulation, etc. That is the reason why, for faster patients' recovery, special attention should be paid to the optimal nutrition. Milk fermentation by *Lactobacillus bulgaricus* (*L. bulgaricus*) strains contributes to a faster going-off of the metabolic processes, to get higher quantities of free amino acids, di-, tri-, and polypeptides, and as a consequence, an easier milk protein absorption by human organism.

Table 1. Basic characteristics of the products from Biostim LBS (Biomilk) series.

INGREDIENTS /100g dry product	BIOSTIM LBS
Water	6%
Dry ingredients	94%
Total proteins	23-27%
Soluble proteins	7-9%
Fats	16-20%
Total carbohydrates	52-57%
incl. lactose	0-4%
Galactose	13-15.5%
Glucose	13-15.5%
Saccharose	13-16%
Energy value	450 kcal
Pectin	0.9%
Minerals	< 6%
Calcium	1000 mg
Potassium	> 1500 mg
Iron	0.8 mg
Phosphorus	780 mg
Vitamin A	0.20 mg
Vitamin D	122 mg
Vitamin B1	0.23 mg
Vitamin B2	1.54 mg
Vitamin C	2.0 mg
Carotene	0.25 mg
Viable LBB cells	> 1010

One of the basic contemporary components reflecting the correct approach to the patients with intoxication is the maintenance of the protein balance and energy needs. In the probiotic Biostim LBS, the proportion of proteins:fats:carbohydrates is 1:0.7:1.8. This is in accordance with WHO recommendations (i. e., 1:1:2).

In our study, the patients with moderate intoxications were given 125 g Biostim LBS daily, which provided 565 kcal/day, while those with severe intoxications received 250 g Biostim LBS daily, which provided 1130 kcal/day. In the patients from both groups, we observed an improved absorption of the nutrients via enteral route. This is evident from the increased energy intake via that route during the hospital stay. Biostim LBS provided as a mean oral energy intake of 45.36% in the group of patients with moderate intoxication and of 70.86% in the group with severe intoxication.

These results were achieved owing to the correct choice of therapy that included mutually additive enteral and parenteral medications. In none of the patients adverse effects were observed at all concerning the parenteral medications and the enterally introduced Biostim LBS as well. They exerted a beneficial effect on the metabolism, improved the energy levels of the body and contributed to the absorption of the energy and the proteins.

A special attention should be paid to the probiotic properties of the viable cells of *L. bulgaricus*, which favorably influence on the organism of the patients by improvement and regulation of the internal bacterial equilibrium (3,14). Their efficacy in increasing the resistance to internal pathogenic flora, in modulating the immune system, in protecting the antigen translocation in the blood flow, as well as their favourable influence on the hepatic encephalopathy is well known (13-15).

CONCLUSION

In conclusion, it should be emphasized that the correct choice of enteral and parenteral therapy ensures the maximally beneficial effect of the probiotic Biostim LBS for the patients with intoxications.

REFERENCES

1. Àěâêñàí ä ðî â Í . Ĭ ðî áéî òèöèòâ-âî áðèòâ áàèòâðèè í à áúëääðñèî òî èèñâçî ì ëýëî . Ñî òèý, Motion publishing, 2003.
2. Delcenserie V, Martel D, Lamoureux M, Amiot J, Boutin Y, Roy D. Immunomodulatory effects of probiotics in the intestinal tract. *Curr Issues Mol Biol.* 2008;**10**(1-2):37-54
3. Donaldson M S. Nutrition and cancer: A review of the evidence for an anti-cancer diet, *Nutr J.* 2004; **3**: 19.
4. Doron S, Gorbach SL. Probiotics: their role in the treatment and prevention of disease. *Expert Review of Anti-Infective Therapy.* 2006; **4**(2):261-275.
5. Elizabeth C. Verna, Use of probiotics in gastrointestinal disorders: what to recommend? *Therap Adv Gastroenterol.* 2010 September; **3**(5): 307-319.
6. Ezendam J, van Loveren H. Probiotics: immunomodulation and evaluation of safety and efficacy. *Nutrition Reviews.* 2006;**64**(1):1-14.
7. Gill H, Prasad J. Probiotics, immunomodulation, and health benefits. *Adv Exp Med Biol.* 2008; **606**:423-54
8. Gill HS, Guarner F. Probiotics and human health: a clinical perspective. *Postgraduate Medical Journal.* 2004; **80**(947):516-526.
9. Goldin BR, Gorbach SL. Clinical indications for probiotics: an overview. *Clin Infect Dis.* 2008 Feb **1**;46 Suppl 2:S96-100; discussion S144-51.
10. Guidance for substantiating the evidence for beneficial effects of probiotics: prevention and management of infections by probiotics.
11. Huebner ES, Surawicz CM. Probiotics in the prevention and treatment of gastrointestinal infections. *Gastroenterology Clinics of North America.* 2006;**35**(2):355-365.
12. Liu JE, Zhang Y, Zhang J, Dong PL, Chen M, Duan ZP. Probiotic yogurt effects on intestinal flora of patients with chronic liver disease. *Nurs Res.* 2010 Nov-Dec;**59**(6):426-432.
13. Minocha A. Probiotics for preventive health. *Nutr Clin Pract.* 2009 Apr-May; **24**(2):227-41.
14. Nova E, Viadel B, Warnberg J, Carreres JE, Marcos A. Beneficial effects of a synbiotic supplement on self-perceived gastrointestinal well-being and immunoinflammatory status of healthy adults. *J Med Food.* 2011 Jan-Feb;**14**(1-2):79-
15. Qian B, Xing M, Cui L, Deng Y, Xu Y, Huang M, Zhang S. Antioxidant, antihypertensive, and immunomodulatory activities of peptide fractions from fermented skim milk with *Lactobacillus delbrueckii* ssp. *bulgaricus* LB340. *J Dairy Res.* 2011 Feb;**78**(1):72-9.
16. Wolvers D, Antoine JM, Myllyluoma E, Schrezenmeir J, Szajewska H, Rijkers GT. Guidance for substantiating the evidence for beneficial effects of probiotics: prevention and management of infections by probiotics. *J Nutr.* 2010 Mar;**140**(3):698S-712S. Epub 2010 Jan 27.